


Copernicus Service Evolution: toward global scale mapping of surface water dynamic at high resolution

M. Dowell

On behalf of

J.-F. Pekel, A. Cottam, A. Belward

European Commission - Joint Research Centre




Objectives



Addressing some of the key questions related to the surface water dynamics


- Provide relevant data at actionable scales
- Where has surface water occurred over the past 3 decades ?
- When do water bodies fill and empty ?
- What about their inter and intra-annual variability ?
- How likely is it to find water in any given place and month ?
- When and where have new/ex water-bodies formed/disappeared ?
- What form did changes take, in terms of seasonality and persistence ?
- What about trends ?

High variability of water spectral signatures



L8 color composition: Swir2, Nir, Red

Ample opportunities for misclassification





Pixel based classifier



Each pixel of the 3,066,102 Landsat scenes was classified as water, land or non-valid observation

- Expert system classifier
- Evidential reasoning and visual analytics approach
- Uses temporal trajectory of pixels in the multispectral feature space
- Hue/Saturation/Value colour model
- Calibrated based on a large spectral library (64,254 pixels)





**Each pixel of the 3,066,102 Landsat scenes (1.8 PB of data)
was classified as water, land or non-valid observation**

Processing using one CPU would have taken 1,212 years


Processing in Google Earth Engine took 45 days

Water history



The When and Where
of the water presence across 32 years

The stack of 32 years of classified Landsat scenes
constitute the water history




Spatio-Temporal Validation

Based on 40.124 validation samples



Omission < 5%
Commission < 1%



	Landsat 5			Landsat 7			Landsat 8		
	Overall	Seasonal	Permanent	Overall	Seasonal	Permanent	Overall	Seasonal	Permanent
Commission accuracy	99.45%	98.80%	99.56%	99.35%	98.38%	99.50%	99.54%	98.53%	99.66%
Omission accuracy	97.01%	74.91%	98.79%	95.79%	73.82%	97.72%	96.25%	77.40%	99.10%

Thematic Products



The validated water history was used to produce thematic products that document different facets of the surface water dynamics

Maps & Temporal Profiles


- Occurrence
- Occurrence Change Intensity
- Seasonality
- Recurrence
- Water Transition
- Max Water Extent

Full monthly water history

(+Metadata layers)




<https://global-surface-water.appspot.com/>



Sacramento river

Flood irrigation
(mainly rice paddies)






Water Occurrence

Provides information concerning the overall water dynamics.

It's a measurement of the water presence frequency



2 km




Water Occurrence Change Intensity

Documents the direction of change
and its intensity (%)

Decrease

No Change


Increase



Intra-annual variability

Separating
Permanent and Seasonal water bodies,
and assessing the degree of seasonality







Search for a location...

Pixel Coordinates: Lat: 39.584261, Long: -121.452329

Monthly Water Recurrence




Lake Oroville Storage Levels



What form does any change in surface water resource take?


Unchanging and changing water surfaces, and their corresponding seasonality



Search for a location...


Pixel Coordinates: Lat: 39.584261, Long: -121.452329

Monthly Water Recurrence




Water History


Click a bar on the graph to see full monthly history for that year



Monthly Water Presence (2012)




- Permanent ☒
- New Permanent
- Lost Permanent
- Seasonal
- New Seasonal
- Lost Seasonal
- Seasonal to Permanent
- Permanent to Seasonal
- Ephemeral Permanent
- Ephemeral Seasonal




2 km

Lakes on the Tibetan Plateau have **increased in area by 20%** with respect to the 1980s: Grazing land is lost and transport links threatened



The Aral Sea has **lost around 1200 km² per year** since 1986

Some recovery is apparent after 2015




High-resolution mapping of global surface water and its long-term changes

Jean-François Pekel¹, Andrew Cottam¹, Noel Gorelick² & Alan S. Belward¹

The location and persistence of surface water (inland and coastal) is both affected by climate and human activity¹ and affects climate^{2,3}, biological diversity⁴ and human wellbeing^{5,6}. Global data sets documenting surface water location and seasonality have been produced from inventories and national descriptions⁷, statistical extrapolation of regional data⁸ and satellite imagery^{9–12}, but measuring long-term changes at high resolution remains a challenge. Here, using three million Landsat satellite images¹³, we quantify changes in global surface water over the past 32 years at 30-metre resolution. We record the months and years when water was present, where occurrence changed and what form changes took in terms of seasonality and persistence. Between 1984 and 2015 permanent surface water has disappeared from an area of almost 90,000 square kilometres, roughly equivalent to that of Lake Superior, though new permanent bodies of surface water covering 184,000 square kilometres have formed elsewhere. All continental regions show a net increase in permanent water, except Oceania, which has a fractional (one per cent) net loss. Much of the increase is

from reservoir filling, although climate change¹⁴ is also implicated. Loss is more geographically concentrated than gain. Over 70 per cent of global net permanent water loss occurred in the Middle East and Central Asia, linked to drought and human actions including river diversion or damming and unregulated withdrawal^{15,16}. Losses in Australia¹⁷ and the USA¹⁸ linked to long-term droughts are also evident. This globally consistent, validated data set shows that impacts of climate change and climate oscillations on surface water occurrence can be measured and that evidence can be gathered to show how surface water is altered by human activities. We anticipate that this freely available data will improve the modelling of surface forcing, provide evidence of state and change in wetland ecotones (the transition areas between biomes), and inform water-management decision-making.

Between any two points in time, part of the Earth's surface is constantly underwater and part is never underwater, with the remainder fluctuating between these extremes. Coastlines and lake and river boundaries advance and retreat, rivers meander, new permanent lakes form and

**DOI: 10.1038/nature20584**<https://global-surface-water.appspot.com/>



Joint Research Centre
Global Surface Water

Data Access

License

All data here is produced under the Copernicus Programme and is provided free of charge, without restriction of use. For the full license information see the [Copernicus Regulation](#).

Publications, models and data products that make use of these datasets must include proper acknowledgement, including citing datasets and the journal article as in the following citation

Citation

Jean-François Pekel, Andrew Cottam, Noel Gorelick, Alan S. Belward, High-resolution mapping of global surface water and its long-term changes. Nature 540, 418–422 (2016). (doi:10.1038/nature20584)

If you are using the data as a layer in a published map, please include the following attribution text: 'Source: EC JRC/Google'

Data Users Guide

For a description of all of the datasets and details on how to use the data please see the [Data Users Guide](#).

Delivery Mechanisms

All of the datasets that comprise the Global Surface Water 1984–2015 are being made freely available using the following delivery mechanisms: Global Surface Water Explorer, Data Doi